

My Asthma Plan



Your asthma plan tells you when to take your asthma medicines.

And what to do when your asthma gets worse.



Name:

1 My daily asthma medicines

 My preventer inhaler is called 	
and its colour is	

- I take _____puff/s of my preventerinhaler in the morning and _____puff/s at night. I do this every day even if I feel well.
- Other asthma medicines I take every day:

My reliever inhaler is called	
and its colour is	

I take____puff/s of my reliever inhaler (usually blue) when I wheeze or cough, my chest hurts or it's hard to breathe.

My best peak flow is

Does doing sport make it hard to breathe?



When my asthma gets worse

I'll know my asthma is getting worse if:

- I wheeze or cough, my chest hurts or it's hard to breathe, or
- I'm waking up at night because of my asthma, or
- I'm taking my reliever inhaler (usually blue) more than three times a week, or
- My peak flow is less than

If my asthma gets worse, I should:

Keep taking my preventer medicines as normal.

And also take_____puff/s of my blue reliever inhaler every four hours.



If I'm not getting any better doing this I should see my doctor or asthma nurse today.

CONSENT FOR EMERGENCY INHALER:

Please sign if you consent for the emergency inhaler in school to be used should it ever be needed:

Date:

Remember to use my inhaler with a spacer (if I have one)



My Asthma Plan

3 When I have an asthma attack

I'm having an asthma attack if:

- My blue reliever inhaler isn't helping, or
- I can't talk or walk easily, or
- I'm breathing hard and fast, or
- I'm coughing or wheezing a lot, or
- My peak flow is less than

When I have an asthma attack, I should:

Sit up — don't lie down. Try to be calm.

Take one puff of my reliever inhaler **every 30 to 60 seconds** up to a total of 10 puffs.

Even if I start to feel better, I don't want this to happen again, so I need tosee my doctor or asthma nurse today.

If I still don't feel better and I've taken ten puffs, I need to call 999 straight away. If I am waiting longer than 15 minutes for an ambulance I should take another_____puff/s of my blue reliever inhaler every 30 to 60 seconds (up to 10 puffs).

My asthma triggers:

Write down things that make your asthma worse

	need	to	see	my	asthma	nurse
E	every	six	mor	nths		

Date I got my asthma plan:

Date of my next asthma review:

Doctor/asthma nurse contact details:



Make sure you have your reliever inhaler (usually blue) with you. You might need it if you come into contact with things that make your asthma worse.

Parents – get the most from your child's action plan

Make it easy for you and your family to find it when you need it

- Take a photo and keep it on your mobile (and your child's mobile if they have one)
- Stick a copy on your fridge door
- **Share** your child's action plan with school, grandparents and babysitter (a printout or a photo).

You and your parents can get your questions answered:

Call our friendly expert nurses

© 0300 222 5800

(9am - 5pm; Mon - Fri)

Get information, tips and ideas

www.asthma.org.uk





Health & care information you can trust